

## Mecanismos, enfoques diagnósticos y estrategias terapéuticas en el paro cardíaco

Ali Abdulridha Obaid<sup>1</sup>, Shahad Abdullah Shwan<sup>2</sup> and Priyanshu Verma<sup>3,\*</sup>

### Resumen

El paro cardíaco es un evento crítico y potencialmente mortal, definido por el cese brusco de la actividad cardíaca efectiva, lo que provoca una interrupción inmediata del flujo sanguíneo hacia los órganos vitales. Sigue siendo una de las principales causas de muerte a nivel mundial y exige una intervención médica inmediata para aumentar las probabilidades de supervivencia y de recuperación neurológica. Las causas del paro cardíaco son diversas e incluyen arritmias ventriculares, infarto agudo de miocardio y cardiopatías estructurales subyacentes. Esta revisión analiza los mecanismos fisiopatológicos que subyacen al paro cardíaco, los principales enfoques diagnósticos utilizados para identificar su etiología y orientar el tratamiento, así como las estrategias actuales de manejo. Se hace especial hincapié en las prácticas de reanimación basadas en la evidencia, en la atención integral posparo cardíaco y en las terapias emergentes dirigidas a mejorar los resultados clínicos de los pacientes afectados.

Insuf Card 2024;19 (1): 13–19.

**Palabras clave:** Atención posparo cardíaco, Reanimación, Paro cardíaco Ventricular arrhythmias, Arritmias ventriculares, Arritmias ventriculares

### Summary

#### *Mechanisms, Diagnostic Approaches, and Treatment Strategies in Cardiac Arrest*

Cardiac arrest is a critical, life-threatening event defined by the abrupt cessation of effective cardiac activity, resulting in an immediate interruption of blood flow to vital organs. It remains one of the leading causes of death globally and demands prompt medical intervention to enhance the chances of survival and neurological recovery. The causes of cardiac arrest are varied and include ventricular arrhythmias, acute myocardial infarction, and underlying structural heart disease. This review examines the mechanisms underlying cardiac arrest, the principal diagnostic approaches used to identify its etiology and guide therapy, and current management strategies. Particular emphasis is placed on evidence-based resuscitation practices, comprehensive post–cardiac arrest care, and emerging therapeutic interventions designed to improve clinical outcomes in affected patients.

**Keywords:** Cardiac arrest, Resuscitation, Post–cardiac arrest care, Ventricular arrhythmias

<sup>1</sup>Mazaya University College, Iraq

<sup>2</sup>Medical Laboratories Techniques, Al-Farahidi University, Iraq Baghdad

<sup>3</sup>Centre for Research Impact & Outcome, Institute of Engineering and Technology, Chitkara University, Rajpura, 140401, Punjab, India

**Correspondencia:** Priyanshu Verma. Centre for Research Impact & Outcome, Institute of Engineering and Technology, Chitkara University, Rajpura, 140401, Punjab, India  
E-mail: priyanshu\_verma1@outlook.in.

Recibido: 22/09/2024

Aceptado: 30/12/2024

## Introduction to Cardiac Arrest

Cardiac arrest is a sudden and often unexpected cessation of heart function, characterized by the abrupt loss of effective circulation and the absence of a palpable pulse. This medical emergency is distinct from a heart attack (myocardial infarction), as it involves the failure of the heart's electrical system rather than a blockage in blood flow [1]. The condition is a leading cause of death worldwide, with survival rates heavily dependent on timely intervention. Without immediate resuscitation efforts, such as cardiopulmonary resuscitation (CPR) and defibrillation, irreversible brain damage or death can occur within minutes [2]. Cardiac arrest can occur in both hospital and out-of-hospital settings, with out-of-hospital cardiac arrest (OHCA) being particularly challenging due to delays in accessing advanced care [3].

The causes of cardiac arrest are diverse and encompass both cardiac and non-cardiac origins. Cardiac causes are primarily arrhythmias, with ventricular fibrillation (VF) and ventricular tachycardia (VT) being the most common. These conditions often result from underlying disorders such as coronary artery disease, cardiomyopathy, or congenital heart defects [4]. Non-cardiac triggers include hypoxia, electrolyte imbalances, drug overdoses, and trauma, all of which can disrupt the heart's electrical or mechanical function [5]. Understanding the multifactorial etiology of cardiac arrest is crucial for developing effective prevention and intervention strategies.

Pathophysiologically, cardiac arrest disrupts the heart's ability to pump blood to vital organs by impairing its electrical system. This results in a rapid decline in oxygen delivery, cellular damage, and multi-organ failure if not promptly reversed [6]. Patients experiencing cardiac arrest often present with sudden collapse, absence of breathing or gasping respirations, and no detectable pulse or blood pressure [7]. The "Chain of Survival" highlights critical steps for improving outcomes in such emergencies, including early recognition and activation of emergency response systems, immediate initiation of high-quality CPR, rapid defibrillation when indicated, and advanced life support and post-resuscitation care [8].

Despite advancements in resuscitation science, survival rates for cardiac arrest remain low, and outcomes are heavily influenced by the timing and quality of intervention. Neurological injuries are a primary concern for survivors, with many experiencing varying degrees of cognitive and functional impairment [9]. Raising awareness about the importance of CPR training, increasing public access to automated

external defibrillators (AEDs), and enhancing emergency medical services are crucial strategies for reducing mortality and improving outcomes [10].

Cardiac arrest arises from a complex interplay of pathophysiological mechanisms. Cardiac arrhythmias, particularly VF and VT, are the most common causes, resulting from abnormalities in the heart's electrical conduction system. VF is characterized by chaotic and uncoordinated electrical activity in the ventricles, rendering the heart unable to generate effective contractions [11]. Sustained VT, if untreated, can degenerate into VF and often occurs in patients with structural heart diseases such as myocardial infarction or cardiomyopathy [12]. Bradycardias and asystole, characterized by severe slowing or complete cessation of electrical activity, are less common but more challenging to treat, with significantly lower survival rates [13].

Ischemia plays a critical role in cardiac arrest by reducing coronary perfusion and triggering electrical instability. Acute blockages in the coronary arteries deprive myocardial tissue of oxygen, impairing the heart's ability to function properly and increasing the likelihood of arrhythmias [14, 15]. Systemic hypoxia and hypoperfusion further exacerbate the condition by depleting cellular energy reserves and causing acid-base imbalances, which impair myocardial contractility and electrical stability [16, 17]. Other contributing factors include electrolyte imbalances and toxicological causes, such as drug overdoses, which can disrupt myocardial excitability and conduction [18].

From a public health perspective, cardiac arrest represents a significant global burden. Efforts to reduce its impact must focus on improving awareness, education, and accessibility to life-saving interventions. Public education campaigns emphasizing CPR training and AED use, along with initiatives to enhance emergency medical services, are essential steps toward reducing the mortality associated with cardiac arrest [10]. By addressing these challenges through a multidisciplinary approach, significant strides can be made in improving survival rates and neurological outcomes.

In conclusion, cardiac arrest is a critical medical emergency with a multifaceted etiology and pathophysiology. Its management requires a co-ordinated response encompassing early recognition, rapid intervention, and comprehensive post-resuscitation care. Continued research and innovation are needed to optimize treatment strategies and mitigate the global burden of this condition.

## Advances in Diagnostic Techniques for Cardiac Arrest

Recent developments in diagnostic tools have significantly enhanced the ability to detect and manage cardiac arrest. These advancements include innovations in electrocardiography (ECG) and advanced medical imaging technologies, enabling early diagnosis, improved risk stratification, and tailored treatment approaches [19].

ECG remains a cornerstone in diagnosing cardiac arrest and its underlying causes. Post-resuscitation ECG is particularly valuable in identifying ischemic changes, such as ST-segment elevation, which can indicate acute myocardial infarction as the underlying cause [20]. High-resolution ECG, including technologies like signal-averaged ECG, can detect subtle electrical abnormalities like late potentials, which are predictive of ventricular arrhythmias [21]. Continuous ECG monitoring through portable and wearable devices has further revolutionized patient care, allowing real-time monitoring of high-risk individuals and aiding in the early detection of arrhythmic events that may precede cardiac arrest [22].

Imaging technologies have also become essential for assessing structural and functional abnormalities contributing to cardiac arrest. Cardiac magnetic resonance imaging (CMR) provides detailed insights into myocardial structure, including scar tissue and fibrosis, which are key markers of arrhythmic risk. It is particularly useful in identifying cardiomyopathies such as hypertrophic cardiomyopathy and arrhythmogenic right ventricular cardiomyopathy [23]. Echocardiography, particularly point-of-care ultrasonography (POCUS), is a rapid and non-invasive tool for evaluating cardiac function during resuscitation and can help differentiate between reversible and irreversible causes of cardiac arrest [24]. Coronary CT angiography is increasingly employed to detect coronary artery obstructions and other structural abnormalities, aiding in identifying ischemic triggers of cardiac arrest [25]. Additionally, nuclear imaging modalities such as PET and SPECT provide valuable insights into myocardial perfusion, viability, and inflammatory conditions associated with cardiac arrest [26].

While not a direct imaging or electrical diagnostic tool, biomarker-based diagnostics have further refined the diagnostic process for cardiac arrest. Biomarkers like troponins and brain natriuretic peptide (BNP) are increasingly integrated with imaging and ECG findings to improve diagnostic accuracy and risk stratification [27].

## Management of Cardiac Arrest: Immediate Interventions

Prompt and effective management of cardiac arrest is critical for survival, as irreversible damage can occur within minutes without adequate circulation and oxygenation. Immediate interventions such as cardiopulmonary resuscitation (CPR) and defibrillation play a pivotal role in restoring cardiac function and preventing long-term complications [28].

The first step in managing cardiac arrest is early recognition and activation of the emergency response system. Recognizing the signs of cardiac arrest, including unresponsiveness, absence of breathing, and lack of a pulse, is vital to initiating timely interventions [29]. Once cardiac arrest is identified, calling emergency medical services (EMS) ensures professional intervention and advanced care [30].

Cardiopulmonary resuscitation (CPR) is the cornerstone of cardiac arrest management. High-quality chest compressions should be performed at a rate of 100–120 per minute and a depth of at least 2 inches (5 cm) in adults, allowing full chest recoil between compressions to maximize venous return [31]. If trained, rescuers should provide rescue breaths at a ratio of 30 compressions to 2 breaths. For those untrained in CPR, continuous chest compressions are recommended until professional help arrives [32]. Interruptions in chest compressions should be minimized to maintain adequate blood flow to vital organs [33].

Defibrillation is another critical component in the management of cardiac arrest. Automated external defibrillators (AEDs) should be used as soon as they become available. Following the device's voice prompts, the heart rhythm is assessed, and a shock is delivered if indicated [34]. Early defibrillation is especially important for ventricular fibrillation (VF) and pulseless ventricular tachycardia (VT), as these rhythms are most amenable to electrical therapy. For healthcare professionals, manual defibrillation allows for tailored shock delivery based on rhythm analysis [35].

Advanced life support (ALS) is often required to stabilize the patient during cardiac arrest management. Airway management, including advanced techniques such as endotracheal intubation, ensures effective oxygen delivery and ventilation. Medications such as epinephrine and amiodarone are administered to address arrhythmias and support circulation [36]. These advanced interventions, combined with high-quality CPR and timely defibrillation, form a comprehensive approach to improving outcomes in cardiac arrest.

## Long-term Management and Rehabilitation

Following survival from cardiac arrest, long-term management and rehabilitation are crucial to prevent recurrence, address underlying conditions, and improve the patient's quality of life. These strategies involve medical, device-based, and rehabilitative approaches tailored to individual patient needs [37]. Implantable cardioverter defibrillators (ICDs) are recommended for patients at high risk of recurrent cardiac arrest due to arrhythmias such as ventricular tachycardia or fibrillation. ICDs continuously monitor heart rhythms and deliver shocks or pacing to restore normal rhythm if necessary [38]. Patients with structural heart diseases, inherited arrhythmias, or a history of sudden cardiac arrest are prime candidates for ICD therapy.

Cardiac resynchronization therapy (CRT) is another important intervention for patients with heart failure and ventricular dyssynchrony, particularly in those with reduced ejection fraction. CRT improves cardiac efficiency and reduces symptoms, thereby decreasing the risk of future cardiac events [39]. Alongside device-based interventions, medical management plays a critical role in long-term care. Pharmacological interventions, including beta-blockers, ACE inhibitors, ARBs, and anti-arrhythmic drugs, are prescribed to manage underlying conditions such as heart failure, hypertension, or arrhythmias. Controlling risk factors like diabetes, hypertension, and hyperlipidemia is equally essential for reducing the overall risk of cardiac events.

Cardiac rehabilitation is an integral part of recovery, combining supervised exercise, education, and psychological support to help patients regain physical strength, manage stress, and adhere to lifestyle changes [40]. Survivors of cardiac arrest often experience anxiety, depression, or post-traumatic stress disorder, making psychological counseling and support groups a vital component of their recovery process. Adopting a heart-healthy lifestyle, including a balanced diet, regular physical activity, smoking cessation, and stress management, is critical for long-term cardiac health.

## Emerging Therapies and Future Directions

The field of cardiac arrest management is witnessing rapid advancements with innovative therapeutic approaches such as regenerative medicine and gene therapy. These cutting-edge technologies hold great promise for improving outcomes and addressing the underlying causes of cardiac conditions that lead to

arrest [42]. Regenerative medicine approaches, including stem cell therapy, aim to restore cardiac function by replacing necrotic or scarred myocardium with new, functional tissue [43]. Exosome-based therapies, derived from stem cells, contain growth factors and microRNAs that promote tissue repair and reduce inflammation, offering a non-cell-based alternative for cardiac regeneration. Advances in bioengineering have enabled the creation of bio-artificial heart tissues and scaffolds, which could repair or replace damaged myocardium in patients with end-stage heart failure or severe post-arrest cardiac dysfunction [44].

Gene therapy is another promising area, with tools like CRISPR/Cas9 being developed to correct hereditary mutations that predispose individuals to arrhythmias and sudden cardiac arrest. Early-stage research focuses on conditions such as long QT syndrome and hypertrophic cardiomyopathy. Adeno-associated viral (AAV) vectors are being explored to deliver therapeutic genes that enhance myocardial function, suppress arrhythmic triggers, or promote angiogenesis in ischemic tissues [45]. Personalized medicine and artificial intelligence (AI) are also playing a transformative role in cardiac care. By combining genetic, molecular, and clinical data, precision medicine enables the development of tailored therapies specific to an individual's risk factors and disease profile. Machine learning algorithms provide AI-driven insights by analyzing large datasets, predicting cardiac arrest risk, and optimizing therapy selection based on patient-specific characteristics.

Novel pharmacological agents targeting specific pathways, such as mitochondrial dysfunction or oxidative stress, are under investigation to improve post-arrest myocardial recovery. Biologic therapies, including cytokine inhibitors and immunomodulators, are being studied to manage inflammatory responses associated with cardiac arrest and resuscitation injury. These emerging therapies and innovative technologies represent a new frontier in the management of cardiac arrest, offering hope for improved outcomes and reduced mortality.

## Role of Public Awareness and Training in Cardiac Arrest Survival

Public awareness and community training programs play a vital role in improving survival rates for cardiac arrest. Since the majority of out-of-hospital cardiac arrests (OHCA) occur in public or residential settings, immediate bystander intervention is often the determining factor in patient survival. Educating

the public about recognizing cardiac arrest, performing CPR, and using automated external defibrillators (AEDs) is critical [46]. Campaigns aimed at educating individuals to identify symptoms such as unresponsiveness and abnormal breathing ensure timely activation of emergency medical services (EMS), reducing delays in initiating life-saving interventions [47]. In addition, awareness programs highlight the importance of locating and using AEDs in public spaces such as airports, schools, and sports facilities. Empowering individuals with the knowledge to operate AEDs confidently during emergencies is a crucial aspect of these initiatives.

Community training programs further enhance public readiness to respond to cardiac emergencies. Hands-only CPR courses teach participants to perform effective chest compressions, enabling immediate response in critical situations. Studies have shown that bystander CPR can double or even triple survival rates [48]. Many workshops also include demonstrations on AED usage, ensuring participants understand how to operate these devices safely and effectively. To maximize reach and impact, training programs often target densely populated areas or regions with low EMS response times, promoting wider dissemination of life-saving skills.

The integration of technology has revolutionized the way public training and awareness are conducted. Mobile applications, such as "PulsePoint," alert trained bystanders to nearby cardiac arrest incidents, facilitating faster intervention. Online learning resources, including e-learning modules and virtual simulations, have made CPR and AED training more accessible to broader audiences, ensuring that more individuals can acquire these critical skills at their convenience.

Encouraging a culture of action is essential for enhancing community responses to cardiac arrest. Public education initiatives aim to reduce fear of intervention by addressing concerns about legal liability or performing CPR incorrectly, thereby empowering individuals to act decisively during emergencies. Programs introduced in schools and workplaces ensure that the next generation is equipped with life-saving skills, fostering a community-wide preparedness for cardiac emergencies. Together, these efforts underscore the importance of public awareness and training in improving survival rates and outcomes for cardiac arrest victims.

## Conclusion

Cardiac arrest remains a critical global health challenge, requiring a comprehensive understanding of its mechanisms, risk factors, and management strategies to improve survival rates and patient outcomes. Advances in diagnostic tools, immediate interventions, and long-term management have significantly contributed to addressing the complexities of this life-threatening condition. Early recognition and timely intervention, including high-quality cardiopulmonary resuscitation (CPR) and defibrillation, are the cornerstones of acute cardiac arrest management. The integration of innovative diagnostic techniques, such as advanced imaging and genetic screening, enables more precise identification of underlying causes, allowing for tailored treatments. Long-term strategies, including implantable cardioverter defibrillators (ICDs), cardiac resynchronization therapy (CRT), and structured rehabilitation programs, enhance quality of life and reduce the risk of recurrence. Emerging therapies like regenerative medicine and gene therapy represent the future of cardiac care, offering the potential to repair damaged myocardium and address the genetic basis of arrhythmic conditions. Additionally, public awareness and community training programs play a pivotal role in improving survival rates, emphasizing the importance of early bystander response and access to automated external defibrillators (AEDs).

## Referencias bibliográficas

- [1] Pura, Agnieszka, Marta Wilk, and Maria Olszowska. "Prognostic value of inflammatory markers in acute coronary syndrome in a population with premature cardiovascular disease (RCD code: VIII)." *Journal of Rare Cardiovascular Diseases* 4.2 (2019): 37-41.
- [2] Raja, S., et al. "Unlocking the potential of polymer 3D printed electronics: Challenges and solutions." *Applied Chemical Engineering* (2024): 3877-3877.
- [3] Raja, S., et al. "Surface functionalization of bio-based polymers in FDM: A pathway to enhanced material performance." *Applied Chemical Engineering* 7.3 (2024).
- [4] Saadh, Mohamed J., et al. "Targeting hypoxic and acidic tumor microenvironment by nanoparticles: A review." *Journal of Drug Delivery Science and Technology* (2024): 105660.
- [5] Saadh, Mohamed J., et al. "Performances of nanotubes and nanocages as anodes in Na-ion battery, K-ion battery, and Mg-ion battery." *Ionics* 30.5 (2024): 2657-2664.

[6] Santos, Diana Katherine Campoverde, et al. "Investigation of ORR and OER Mechanisms by Co-and Fe-doped Silicon Nanocages (Si48 and Si60) and Co-and Fe-doped Silicon Nanotubes (SiNT (5, 0) and SiNT (6, 0)) as Acceptable Catalysts." *Silicon* (2024): 1-13.

[7] Neini, O. D. "Enhanced Growth Hormone and IGF-1 Axis Modulation in Chronic Heart Failure: A Multicenter, Double-Blind, Placebo-Controlled Study". *Journal of Rare Cardiovascular Diseases* 4.6. (2022): 122-126.

[8] Subramani, Raja, et al. "Advancements in 3D printing materials: A comparative analysis of performance and applications." *Applied Chemical Engineering* (2024): 3867-3867.

[9] Pura, Agnieszka, Marta Wilk, and Maria Olszowska. "Prognostic value of inflammatory markers in acute coronary syndrome in a population with premature cardiovascular disease (RCD code: VIII)." *Journal of Rare Cardiovascular Diseases* 4.2 (2019): 37-41.

[10] Thabet, Zahraa Fadel, and Heba A. Abd-Alsalam Al-salame. "Evaluation the Effectiveness of Nephelium lappaceum (NL) Peel Aqueous Extract Against Hepatocellular Carcinoma Induced by Thioacetamide (TAA) in Male Albino Rats." *Pure Sciences International Journal of Kerbala* 1.2 (2024).

[11] Prochwiecz, Anna, Elżbieta Szczepanek, and Dorota Krochmalczyk. "Current views on the use of interferons in the treatment of polycythaemia vera (RCD code: VIII)." *Journal of Rare Cardiovascular Diseases* 4.2 (2019): 34-36.

[12] Alex, John. "Atrial Fibrillation Management: The Combined Approach of Left Atrial Appendage Closure and Catheter Ablation." *Journal of Rare Cardiovascular Diseases* 4.8 (2024): 157-162.

[13] Saadh, Mohamed J., et al. "Incorporation anthracene and Cu to NH<sub>2</sub>-Zr-UiO-67 metal-organic framework: Introducing the simultaneous selectivity and efficiency in photocatalytic CO<sub>2</sub> reduction to ethanol." *Journal of Molecular Structure* 1318 (2024): 139329.

[14] Mustafa, Mohammed Ahmed, et al. "A novel fuzzy M-transform technique for sustainable ground water level prediction." *Applied Geomatics* 16.1 (2024): 9-15.

[15] Mustafa, Mahbuba A., and Soran A. Othman. "Effect of adding natural and synthetic antioxidants to broiler drinking water as antistressor on productivity, antioxidant status and hematological traits under heat stress." *Tikrit Journal for Agricultural Sciences* 24.1 (2024): 94-104.

[16] Mustafa, Harith Ahmed, et al. "Isolation and Identification of Some Types of Pathogenic Bacteria from the Prepuce (Foreskin) of Circumcised Children in Samarra City/Iraq." *HIV Nursing* 22.2 (2022): 2776-2780.

[17] Mohammed, Arkan B., Oday K. Hamad, and Thamer A. Khttab. "Effect of zinc oxide nanoparticles in drinking water on growth rate, biochemical parameters, and intestinal histology of broilers." *Advances in Agriculture* 2023.1 (2023): 8523516.

[18] Kadhum, Hussam H., et al. "Potential pharmaceutical applications and molecular docking study for green fabricated ZnO nanoparticles mediated Raphanus sativus: In vitro and in vivo study." *Nanotechnology Reviews* 13.1 (2024): 20240113.

[19] AL-Aameli, Muna Hussain, Rajaa Ali Moheiseen Al-Taee, and H. A. Alsalam. "Histological & physiological alternations in the aorta & heart in relation with cholesterol diet in male albino rat." *Indian J. Forensic Med. Toxicol* 13 (2019): 313-318.

[20] Alabbasy, Raghad Hazem, et al. "Histological study of the effect of some oncology drugs on heart muscle." *AIP Conference Proceedings*. Vol. 2977. No. 1. AIP Publishing, 2023.

[21] Al-Abodi, Hiba Riyadh, et al. "Novel gold nanobiosensor platforms for rapid and inexpensive detection of Vibrio cholerae." *Reviews and Research in Medical Microbiology* 31.2 (2020): 70-74.

[22] Al-Bahrani, Hussein S., et al. "Determining new way to detect hydrocarbon polluted water at vertical and horizontal spaces." *Materials Today: Proceedings* 42 (2021): 2071-2075.

[23] Ali, Raghad Hassan, et al. "Assessment of Serum Cystatin C and Apolipoprotein Levels in Chronic Kidney Disease Patients." *Iraqi Journal of Medical and Health Sciences* 1.1 (2024): 8-13.

[24] AL-qanbar, Mohammed Majid, Wefak Jbori AL-Bazi, and Hepa A. Abd-Alsalam. "The effect of Hyperhomocysteinemia on the Osteoclasts activity in Male New Zealand White Rabbits." *Research Journal of Pharmacy and Technology* 15.12 (2022): 5443-5448.

[25] Al-Salam, Yasser, Mohammed J. Mohammed, and Hani Sabbar Ayed. "Effect of Molecular Substitution of Oats and Chia Powder for Animal Fats on Chemical Indicators and Physical Properties in the Manufacture of Freeze-Preserved Beef Burgers." *Tikrit Journal for Agricultural Sciences* 24.1 (2024): 170-179.

[26] Al-Omairi, Baida Ajeeal Badir, and Len Nokes. "The Impact of Twisting on The Extent of Lower Lumbar Spine Lateral Flexion Among Healthy Subjects." *Iraqi Journal of Medical and Health Sciences* 1.1 (2024): 22-26.

[27] Alsalam, Heba A. Abd-Alsalam. "Study the Effect of Lycium barbarum Polysaccharide on Bone and Thyroid Gland in Hyperlipidemic Healthy Male

Albino Rats." *Scopus Ijphrd Citation Score* 10.7 (2019): 764.

[28] Alsalame, Heba A. Abd-Alsalam. "Study Effects of Nigella Sativa Seeds Oil in Some Physiological Parameters in Experimental Heart Failure Induced by Ivabradine in Male Rats." *Indian Journal of Forensic Medicine & Toxicology* 14.3 (2020): 2609-2613.

[29] Alsalame, Heba A. Abd-Alsalam, and Layla S. Laylani. "Evolution the effectiveness of Vernonia amygdalina in some physiological parameters against hepatic and renal injury in male rats: Evaluating the Efficacy of Vernonia amygdalina on Physiological Parameters in Ameliorating Hepatic and Renal Injury in Male Rats." *Tikrit Journal for Agricultural Sciences* 24.2 (2024): 298-310.

[30] Alsalame, NHeba A., et al. "Protective Role of Alchoholic Extract of Fennel Seed in Nephrotoxicity Induced by Cisplatin in Male Rabbits." *Biochemical & Cellular Archives* 18 (2018).

[31] Leo Oscar, George Arthur and k. Amelia. "Cardiac Amyloidosis in 2023: Transformative Advances in Diagnosis and Therapy" *Journal of Rare Cardiovascular Diseases* 4. 7 (2023) : 145-150

[32] Ortiz, Daniela Tatiana Castañeda, et al. "Exploring the photovoltaic performance of boron carbide quantum dots doped with heteroatoms: A DFT analysis." *Diamond and Related Materials* 143 (2024): 110933.

[33] Fadhil, K. B., M. A. A. Majeed, and M. A. Mustafa. "Electronic study of fresh enzyme complexes of antifungal drugs-P450 and Aspergillus kojiic acid biosynthesis. W: w saccharose flavus: fructose as a substratum." *Annals of Tropical Medicine and Health* 22 (2019): 65-72.

[34] George, Mike. "A Comprehensive Framework for Rare Disease Management in Primary Care." *Journal of Rare Cardiovascular Diseases* 4.8 (2024): 163-165.

[35] Hama Hasan, Torken Ahmed, et al. "Effect of *Equisetum Arvense* Phenolic Extract in Treatment of *Entamoeba Histolytica* Infection." *Systematic Reviews in Pharmacy* 11.11 (2020).

[36] Hsu, Chou-Yi, et al. "Amphiregulin in infectious diseases: Role, mechanism, and potential therapeutic targets." *Microbial Pathogenesis* 186 (2024): 106463.

[37] Hsu, Chou-Yi, et al. "Exploiting the immune system in hepatic tumor targeting: unleashing the potential of drugs, natural products, and nanoparticles." *Pathology-Research and Practice* (2024): 155266.

[38] Jalal, Bnar J., and Mustafa RM Alqaisi. "Improving the production and quality of white button mushroom (*Agaricus bisporus*) by adding biochar and ash to the casing layer." *Tikrit Journal for Agricultural Sciences* 24.1 (2024): 22-33.

[39] Jawad, Zainab Nizar. "The Association of Urokinase Gene 3'-Utr T/c Polymorphism With Urinary Bladder Cancer." *Biochemical & Cellular Archives* 18.1 (2018).

[40] Jawad, Zainab Nizar. "Molecular Diagnosis of CD14 and MnSOD Genes and their Effect on Asthma in Holy Karbala, Iraq." *Indian Journal of Public Health Research & Development* 10.8 (2019).

[41] Jawad, Zainab Nizar. "Molecular detection of caspase 3, 8, 9 genes and ADIPOR1 (rs2275738) polymorphism in colorectal cancer." *Applied Nanoscience* 13.8 (2023): 5365-5368.

[42] Jawad, Zainab Nizar. "Association between APC gene SNPs and the risk of occurrence of colorectal cancer." *AIP Conference Proceedings*. Vol. 2414. No. 1. AIP Publishing, 2023.

[43] Jawad, Zainab Nizar, and Kamal AR. "Association of Additive Risk of Pioglitazone Use with the Presence of CYP1A1 Polymorphisms in the Occurrence of Bladder Cancer." *Systematic Reviews in Pharmacy* 11.5 (2020).

[44] Jawad, Zainab Nizar, and Weaam Awad. "Association of urokinase and Vitamin D receptor genes SNPs and urolithiasis in an Iraqi population." *Meta Gene* 24 (2020): 100679.

[45] Jawad, Zainab Nizar, and Weaam Awad. "The effect of montelukast on CD14 gene expression in asthmatic patient in Kerbala." *AIP Conference Proceedings*. Vol. 2290. No. 1. AIP Publishing, 2020.

[46] Jawad, Zainab Nizar, Zuhair Mohammed Ali Jeddooa, and Riadh Rasheed Al-Toama. "Association of MDM2 (T309G) gene polymorphism with interstitial urinary bladder cancer." *Biochemical & Cellular Archives* 18.2 (2018).

[47] Jawad, Zainab Nizar, Kamal Abdul Rasool, and Weaam Awad. "The relationship between smoking and Urokinase gene 3'-UTR T/C expression on occurrence of bladder cancer." *Indian Journal of Forensic Medicine & Toxicology* 14.1 (2020): 1169-1171.

[48] Kadham, Shaymaa Maki, et al. "IoT and artificial intelligence-based fuzzy-integral N-transform for sustainable groundwater management." *Applied Geomatics* 16.1 (2024): 1-8.